



BRISK PAIN RELIEF¹

(Piroxicam β-Cyclodextrin) **BRYSK** has shortest Tmax and longer half life providing faster & prolonged analgesia as compared to traditional NSAIDs.

Exceptional GI Safety Vs Other NSAIDs²

GI Events with PBC, Piroxicam and Other Reference Agents in 28 Chronic Treatment Studies from a Pooled Safety Analysis

	Piroxicam- β-Cyclodextrin ^a	Piroxicam	Others ^b
	N=15040	N=579	N=314
ADRs	%	%	%
Abdominal	0.33	5.87	0.96
Constipation	0.33	3.28	0.00
Dyspepsia	0.57	11.92	5.41
Gastritis	0.33	1.90	6.05

^aIncludes all dosage forms and all doses of PBC

^bOther reference agents included tenoxicam, diclofenac/diclofenac sodium, meclofenamate sodium, nabumetone, etodolac, ketorolac, tromethamine and droxicam.

Ref: 1. Adis International Limited, Auckland; New Zealand drugs48(6);907-929 194
2. Current Medicinal Chemistry, 2013, 20, 2415-2437

Indications	Dosage
Osteoarthritis.	OD
Rheumatoid arthritis.	OD
Low back pain.	OD
Sprains and Strains.	OD
Primary dysmenorrhea.	OD
Dental pain.	OD



Full Prescribing Information Available on Request

Dyson Research Laboratories (Pvt) Ltd.

28th-KM Ferozepur Road, Lahore, Pakistan



15



brisk
PAIN RELIEF